

# Goliath® Plus

Gel Bait

## Evolution is a giant plus.

Goliath® Plus Gel Bait sets a new standard. Unique dual active formulation to control not only cockroaches but now ants, and silverfish. It is palatable, cost-effective, convenient, and reduces resistance risk.

- Superior palatability for up to 3 months resulting in increased bait uptake
- Unsurpassed efficacy providing long term results in the toughest situations
- Transfer Effect™ ensuring that the powerful actives provide quicker population control
- Unique dual active advantage, reduces the risk of resistance
- Cost effective gel bait with one cartridge providing up to 1,000 placements



For more information on Goliath Plus Gel Bait, visit [pest-control.basf.com.au](http://pest-control.basf.com.au) or contact your local BASF representative on **1800 558 399**.

 **BASF**  
We create chemistry

# Goliath® Plus

## Gel Bait

Goliath Plus Gel Bait contains 0.4 g/kg of Fipronil and 0.4 g/kg Chlorfenapyr, meaning only small amounts of bait are required for the powerful Goliath Transfer Effect to start working through cockroach, ant and silverfish populations. The active loading and superior formulation not only make Goliath Gel Bait highly effective but also result in the most economical cockroach bait on the market with the added benefit of ants and silverfish on the label. A cost-efficient solution with one cartridge providing up to 1,000 individual bait placements.

Situation	Pest	Rate	Critical Comments
Domestic, commercial, industrial and public buildings and facilities (including but not limited to food processing establishments, shops, factories, farm buildings, hotels and backpacker accommodation, ships, trains, aeroplanes, offices, schools, storerooms, hospitals, barracks, private houses and external areas associated with such establishments).	German and other common cockroaches, ants, silverfish.	Apply 1-3 spots of gel per m <sup>2</sup> . Recommended spot size is 0.03-0.06 g. Increase the number of spots (up to 3) and spot size depending on severity of infestation.  Ants: Apply lines of gel approximately 10-25 mm long for highly active ant foraging trails.	Gel placement should be at or near areas where cockroaches, ants or silverfish gather, along active foraging trails, nest sites or known active areas. Gel may be placed in bait stations or bait cafes, where required.  Inspect bait placements periodically for consumption or attractiveness and reapply as required.

Refer to the product label for full directions and use table.



## Packaging

4 x 30 g cartridges

For more information on Goliath Plus Gel Bait, visit [pest-control.basf.com.au](https://pest-control.basf.com.au) or contact your local BASF representative on **1800 558 399**.

### ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.

This fact sheet is intended as general advice. Disclaimer: The information submitted in this publication is based on current BASF knowledge and experience. In view of the many factors that may affect its application, this data does not relieve the user from carrying out their own tests. The data does not imply assurance of certain properties or of suitability for a specific purpose. It is the responsibility of the user to ensure that any proprietary rights and existing laws and legislation are observed.